

405 EXTENDED TEMPORAL LIFT (add on)

SEE add-on sutures and instruments

almost always done with facelift, but can be done alone

Usually under LOCAL ANESTHESIA;

updated July 27, 2016 PAGE 1

OPERATIVE TIME:	add 1 hour to facelift, or 1 ½ hours alone
ANESTHESIA NOTES	<u>LOCAL ANESTHESIA</u> : patient receives preop Valium, Demerol, and Clonidine. Moderate sedation protocol. Nasal O ₂ cannula with CO ₂ monitor
PATIENT POSITION SETUP	<u>HOLDING AREA</u> : On Stryker eye bed. Start IV on right side. Baer hugger. Apply warm Bovie pad, EKG pads, wrist restraints, etc. in holding. Give IV Kefzol, Zofran, Decadron and ½ to 1 mg Versed. Warm blankets, kisses and hugs. <u>IN OR</u> : Position patient bed flat, pillow under knees, terry cloth towel, gel pad, towels for headrest. [2] blue Chux under head, blankets to tuck arms. Rotate clockwise 135°. Nasal prongs with CO ₂ monitor. - Aquaphor to lips. - Kerlix roll wrist restraints. - Hair ties, comb and shaver box - extra pillow in room for end of case
ROOM SETUP	Warm room to 70°, then lower to 68° after patient. draped Warm up Bovie grounding pad and EKG leads in warmer [for local] [2] ring stands for dirty sponges; [1] Double ring stand for ice bath [3] Mayo stands, [1] large back table Bovie machine at foot of bed; bipolar foot pedal taped to floor
EQUIPMENT	Headlight w/ [2] light sources, Gwen's foot pad [2] white hydraulic pump up stools
PREP	Chlorxylenol (PCMX), 4x4 sponge box - <i>do not add water</i> . Prep lightly to face, eyelids, forehead, neck, hair
DRAPING	Head sheet w/ 2 towels, 1000 drape, split sheet
CAUTERY	✓ Valleylab flat short silver tip (E1450-X)
SUCTION	[2] blue 3/16" suction tubing for Saf-T-Vac & Yankauer
SOLUTIONS IRRIGATION	[3] liters NS irrigation, [2] Asepto syringes, [1] saline bowl COLD SALINE BATH SETUP : get crushed ice in cooler, place ice into large basin on double ring stand, cover with warmer drape, place small sterile basin on top of warmer drape, pour saline in sterile basin
NEEDLES SYRINGES	[4] 3cc syringes, 30g needles ➤ for solution E-2 [2] 10 cc syringes, 1 each 25 and 27g needles ➤ for solution E-2 [2] 16g angiocath [2] 30cc syringes ➤ for Gwen irrigation
LOCAL ANESTHETIC & SOLUTIONS	SOLUTION E-2: Mix 50/50: 20cc 1% Lidocaine w/ epi + 20cc 0.5% Marcaine w/epi
MEDICATIONS (for local)	[4] Versed; [4] Fentanyl; [1] Zofran; [1] Kefzol; [2] Bactroban ointment tubes

SETS TO OPEN

Dr C Facelift & Dr. C Special Plastic set

INSTRUMENTS

■ **FORCEPS**

01's Adsons, 08's

■ **SCISSORS**

Gorney, short ceramics, ceramic Metz, str Metz, str Mayo

■ **NEEDLE HOLDERS**

[2] Mayo Hagar NH
[2] Aesculap serr NH

■ **RETRACTORS**

Breast, Maxwell, Army-Navy, L-shaped retractor

■ **KNIFE | BLADES**

[4] No. 10 blades

■ **OTHER**

STERILE SUPPLIES

- 1000 Steri-drape
- [2] needle counters
- cotton balls for ears
- ½" steri-strips (hold for general)
- plastic drape for iced saline
- [4] surgical marker pens
- Mastisol vial
- magnetic pad
- [4] blue plastic bowels
- Plastic Yankauer
- [3] med glasses

SPONGES

[4] Raytec, [1] Lap

SUTURES

No	Type	SN	status	
	>>> OPEN IN ADDITION TO FACELIFT SUTURES <<<			
[4]	3-0 nylon PS-1	1663	open 4	
[4]	3-0 Chromic PS-2	1638	open 4	
[4]	4-0 Chromic PS-2	1637	open 4	
[4]	0 STRATAFIX CP-2, 26mm	SXPD2B-415	open 2	hold 2
[2]	3-0 STRATAFIX FS, 26mm	SXMD2B-412	open 2	

DRAINS

No drains are used for temporal lift

DRESSINGS

place on Mayo stand
start setup after injections
over

- Knife blade to cut rubber bands
- [2] Bactroban ointment tubes
- tongue blade
- [2] packs 4x4's soft sponges
- [1] Abd pad
- Lister bandage scissors
- [1] 4" Ace
- [2] 3x18 Vaseline gauze
- [2] Kerlix
- [2] Safety pins
- Dr. C sterile gloves

POSTOPERATIVE

Remove head cushion, put 2nd pillow under head

DATES MODIFIED

Jul 2016, Jan 2017