

413 BROWLIFT

set up same as **FACELIFT**.
may include Facelift, Upper and/or Lower Blepharoplasty
may include resection of glabellar muscles - ASK

updated January 4, 2017 Page 1

OPERATIVE TIME	2½ hours UNDER LOCAL if done alone; 6 hours total if done with facelift, usually under GENERAL ANESTHESIA
ANESTHESIA NOTES Note: for general anesthesia use ENT position on regular OR table	<u>LOCAL ANESTHESIA:</u> patient receives preop Valium, Demerol, and Clonidine. Moderate sedation protocol. Nasal O ₂ cannula with CO ₂ monitor <u>GENERAL ANESTHESIA:</u> SCD's, Foley catheter, ✓ regular OR table in ENT position - rotate <i>clockwise</i> after induction ✓ Dr. C to sew in tube with 0-Vicryl CT1, needle holder, scissors ✓ Accordion extension, ✓ straight connector, ✓ long circuit, ✓ regular ET tube
PATIENT POSITION SETUP	<u>HOLDING AREA:</u> On Stryker eye bed. Start IV on right side. Baer hugger. Apply warm Bovie pad, EKG pads, wrist restraints, etc. in holding. Give IV Kefzol, Zofran, Decadron and ½ to 1 mg Versed. Warm blankets, kisses and hugs. <u>IN OR:</u> Position patient bed flat, pillow under knees, terry cloth towel, gel pad, towels for headrest. [2] blue Chux under head, blankets to tuck arms. Rotate clockwise 135°. Nasal prongs with CO ₂ monitor. - Aquaphor to lips. - Kerlix roll wrist restraints. - Hair ties, comb and shaver box - extra pillow in room for end of case
ROOM SETUP	Warm room to 70°, then lower to 68° after patient. draped Warm up Bovie grounding pad and EKG leads in warmer [for local] [2] ring stands for dirty sponges; [1] Double ring stand for ice bath [3] Mayo stands, [1] large back table Bovie machine at foot of bed; bipolar foot pedal taped to floor
EQUIPMENT	Headlight w/ [2] light sources, Gwen's foot pad [2] white hydraulic pump up stools
PREP	Chlorxyleneol (PCMX), 4x4 sponge box - <i>do not add water</i> . Prep lightly to face, eyelids, forehead, neck, hair
DRAPING	Head sheet w/ 2 towels, 1000 drape, split sheet
CAUTERY x 2	Bovie 20/20 pure w/ flat silver tip. Bi-polar with cord, disposable green bipolar forceps set to 15. [2] bovie holsters.
SUCTION x 2	[2] suction tubing: Saf-T-Vac and baby Yankauer
SOLUTIONS IRRIGATION	[3] liters NS irrigation, [2] Asepto syringes, [1] saline bowl COLD SALINE BATH SETUP: get crushed ice in cooler, place ice into large basin on double ring stand, cover with warmer drape, place small sterile basin on top of warmer drape, pour saline in sterile basin.
NEEDLES SYRINGES	NEEDLES: [4] 25g 1½", [4] 27g 1¼", [4] 30g ½" SYRINGES: [4] 10cc syringes, [4] 3cc syringes [2] 25g 1½" needles placed on [2] 10cc syringes ➤ Solution F [2] 27g 1¼" needles placed on [2] 10cc syringes ➤ Solution F [4] 30g ½" needles placed on [4] 3cc syringes ➤ Solution E-2 [2] 16g angiocath [2] 30cc syringes ➤ for Gwen irrigation
LOCAL ANESTHETIC & SOLUTIONS	Solution F: buffered Lidocaine 0.5% w/epi 50cc + NaBicarb 8.4% 5cc Solution E-2: mix 50cc 1% lidocaine w/ epi + 50cc 0.5% Marcaine w/epi
MEDICATIONS	[4] Versed; [4] Fentanyl; [1] Zofran; [1] Kefzol; [2] Bactroban ointment tubes BSS, Surgilube for eyes Decadron 8 mg IV Purelube ophthalmic ointment

3 SETS TO OPEN

[1] Facelift set, [2] Plastic set, [3] bleph fine facial set - *HOLD*
 ->NOTE: ask if doing glabella muscles, if not, *DO NOT OPEN BLEPH SET*

INSTRUMENTS [3] Mayo stands; [2] Basin stands

- 1st MAYO – Opening and dissection
- 2nd MAYO – Local infiltration: syringes and needles, then dressings
- 3rd MAYO – Closure instruments

■ **FORCEPS** 01 - 05 - 08 Adsons, Smooth Adson,
[for glabella add -> Bishop Harmon smooth]

■ **SCISSORS** regular super cut Metz, str Metz, mini-Metz, str mayo, Gorney
[for glabella add -> Stevens tenotomy, Gradles, baby, mama and papa bears]

■ **NEEDLE HOLDERS** [4] Mayo Hagar NH, [4] Aesculap Crile Wood NH

■ **RETRACTORS** Maxwell facelift, narrow Freeman facelift, [2] large facelift, Senn, Chin retractor

■ **KNIFE | BLADES** [6] No. 10 blades; 3 round knife handles

■ **FOR GLABELLA ADD ->** *Bishop Harmon smooth*
Stevens tenotomy, Gradles, baby, mama and papa bears

STERILE SUPPLIES

- 1000 Steri-drape
- [2] needle counters
- cotton balls for ears
- ½" steri-strips (hold for general)
- Disposable bipolar cord #10-4001
- warmer plastic drape for iced saline
- Tegaderm for eyes, cut ½
- [8] surgical marker pens
- Mastisol (hold for general)
- magnetic pad
- [4] blue plastic bowls
- Plastic Yankauer
- [3] med glasses
- Purelube ophthalmic ointment

SPONGES [4] Raytec, [1] Lap

SUTURES

No	Type	SN	status
[4]	3-0 Nylon PS-1	16631	open
[4]	0 STRATAFIX CP-2, 26mm	SXPD2B-415	open
[2]	3-0 STRATAFIX FS, 26mm	SXMD2B-412	open
[4]	3-0 Chromic PS-2	1638	open
[2]	4-0 Chromic	1637	open
[1]	0 Ethibond CT1 - <i>General only</i>		hold

DRAINS NONE, unless doing facelift at same time

DRESSINGS on MAYO

- [1] Bactroban ointment
- tongue blade
- [2] 4x4's
- [1] ABD
- only [1] 18x3 Vaseline gauze
- [2] Kerlix
- 4" Ace
- Lister's bandage scissors

POSTOPERATIVE

Cold compress, gauze soaked in ice saline, for eyes
 Remove head cushion, put pillow under head, lift head of bed

DATES MODIFIED

Jun 2006, Aug 2007, Mar 2008, Dec 2008, May 2009, Mar 2011, Aug 2011, Mar 2012, Jul 2013, Nov 2013, Sep 2014, Aug 2016, Jan 2017