

416 CHIN or CHEEK IMPLANTS

Chin implant is placed through a submental incision.

Cheek implants are placed through an intra-oral incision.

Note: This procedure can be done alone but is usually added to another, larger procedure such as facelift, submental lipectomy or rhinoplasty

Updated March 3, 2013Page 1

OPERATIVE TIME: add ½ hour to main procedure, 1hr if alone

FREQUENCY 1-2 per year

ASSISTANT: Gwen

CPT CODE(S):

ICD9 CODE: V50.1 Elective Cosmetic, 744.9 face Neck

ANESTHESIA NOTES

In holding patient receives oral pre-sedation:
Valium, Demerol, Benadryl, and/or Clonidine.
Nasal O₂ cannula with CO₂ monitor.
If under general, see facelift sheet.

PATIENT POSITION | SETUP

HOLDING AREA:
On Stryker Eye Bed Dr. C marks pt. in holding.
Have holding start IV on right side, O₂ in Holding.
IN OR:
Pillows under knees.
Gel pad, w/ rolled Teri cloth towel.
Kerlix wrist retainers
[2] blue Chux under head, [2] blankets to tuck arms
Nasal prongs with CO₂ monitor.
Aquaphor to lips

TABLE POSITION

rotate clockwise 135°
side rails down on Stryker, patient belt, Kerlix wrist restraints

ROOM SETUP

normal temperature DO NOT pre-warm
Warm up Bovie grounding pad and EKG leads
NO Bair hugger

EQUIPMENT

Headlight and [2] light sources, [2] stools

SPECIALS | IMPLANTS

Dr. C will bring implants and sizers.

PREP

PCMX or Chlorxylenol, lightly with very little H₂O

DRAPING

Head sheet, 1000 drape, split sheet, 4 towels,
[2] rigid bovie holsters, [2] packs green towels
Custom ENT pack #315360

CAUTERY

Bovie 20/20 pure, 3 tips: Flat silver tip, long needle tip (018A),
[2] holsters

SUCTION

[2] suction tubing, Saf-T-Vac, Yankauer, baby Yankauer

SOLUTIONS | IRRIGATION

! NS irrigation
! Small blue bulb syringe.
! Kefzol antibiotic irrigation in separate bowl

NEEDLES | SYRINGES

[2] 25g 1½" needles
[2] 27g 1" needle on [2] 10cc syringe
[2] 30g ½" needles on [2] 3cc syringes

LOCAL ANESTHETIC

Mix 50/50:
1% lidocaine w/ epi 1:100,000 - 20cc, [2] vials
0.5 % marcaine w/epi 1:100,000 - 20cc [2] vials

MEDICATIONS

! [2-4] Versed
! [2-4] Fentanyl
! [1] Zofran
! [1] Kefzol
! Bacitracin ointment 2oz

SETS TO OPEN	Dr. C Facelift, Minor set, and Rhino set for elevators and retractors
INSTRUMENTS USED (ON 2 MAYO STANDS)	
○ FORCEPS	01- 03 - 05 Adsons, Smooth Adson, Adson Brown, Cushing, Short Debaquey, Tissue forceps
○ SCISSORS	cvd and str Stevens, Small Freemans, cvd and str Metzenbaums, Joseph
○ NEEDLE HOLDERS	Webster smooth and serrated, Crile-Wood
○ RETRACTORS	Senn, Ragnell, L-retractor, small facelift, chin senn, Converse nasal.
○ KNIFE BLADES	[2] No 15, [2] No 10, round blue handle needle holder
○ ELEVATORS	Joseph, Freer, Tebbetts
○ OTHER	Cutting board w/silastic sheet

Note: what to pull extra when doing chin or cheek implants with other major procedure	ELEVATORS: Joseph, Key, Freer, Tebbetts (Rhino set) RETRACTORS: Converse nasal, Ragnell, Senn (Rhino set) SCISSORS: Joseph, curved black iris (Facelift set) LOCAL: same as main procedure OTHER: ½" steri-strips
--	--

STERILE SUPPLIES	! steri-drape 1000 ! [2] needle counters ! med glasses ! surgical marker pens	! labels for local ! magnetic pad ! ½" steri-strips ! Mastisol		
SPONGES	Raytec			
NON-STERILE SUPPLIES				
	No	Type	SN	status
SUTURES	M	[2] 3-0 Vicryl SH	J416	hold
	M	[2] 5-0 Monocryl P-3	J493	hold
	M	[2] 4-0 Chromic PS-2	1638	hold
	M	[2] 5-0 Chromic P-3	Y493	hold
	M	[2] 5-0 Plain P3	686	hold
DRAINS	none, unless doing simultaneous facelift			
DRESSINGS	<i>ASK before opening:</i> Bacitracin ointment, Vaseline gauze, Mastisol, ½ inch Steri-strips, 4" Kling or Kerlix, 4" Ace. crushed ice in cooler			
POSTOPERATIVE	Remove head cushion, put 2 nd pillow under head, lift head of bed			
DATES MODIFIED	March 2008, December 2008, October 2011, Dec 2012, Mar 2013			