

419 Excision Buccal Fat Pads (intra-oral approach)

In this operation, buccal fat pads are removed through an intra-oral incision.

Usually under LOCAL ANESTHESIA;

Can be done alone, but often done in conjunction with facelift

updated January 20, 2017 page 1

OPERATIVE TIME: 1½ - 2 hours	
ANESTHESIA NOTES Note: for general anesthesia use ENT position	<u>LOCAL ANESTHESIA:</u> patient receives preop Valium, Demerol, and Clonidine. Moderate sedation protocol. Nasal O ₂ cannula with CO ₂ monitor
	<u>GENERAL ANESTHESIA:</u> SCD's, Foley catheter, ✓ regular OR table in ENT position - rotate <i>clockwise</i> after induction ✓ Dr. C to sew in tube with 0-Vicryl CT1, needle holder, scissors ✓ Accordion extension, ✓ straight connector, ✓ long circuit , ✓ regular ET tube
	PATIENT POSITION SETUP <u>HOLDING AREA:</u> On Stryker eye bed. Start IV on right side. Baer hugger. Apply warm Bovie pad, EKG pads, wrist restraints, etc. in holding. Give IV Kefzol, Zofran, Decadron and ½ to 1 mg Versed. Warm blankets, kisses and hugs. <u>IN OR:</u> Position patient bed flat, pillow under knees, terry cloth towel, gel pad, towels for headrest. [2] blue Chux under head, blankets to tuck arms. Rotate clockwise 135°. Nasal prongs with CO ₂ monitor. - Aquaphor to lips. - Kerlix roll wrist restraints. - Hair ties, comb and shaver box - extra pillow in room for end of case
	ROOM SETUP Warm room to 70°, then lower to 68° after patient. draped Warm up Bovie grounding pad and EKG leads in warmer [for local] [2] ring stands for dirty sponges; [1] Double ring stand for ice bath [3] Mayo stands, [1] large back table Bovie machine at foot of bed; bipolar foot pedal taped to floor
EQUIPMENT	Headlight and extension box, [2] light sources, [2] stools
PREP	Chlorxylenol (PCMX), 4x4 sponge box - <i>do not add water.</i>
DRAPING	Head sheet, 1000 drape, split sheet, 4 towels, [2] rigid bovie holsters, [2] packs green towels Custom ENT pack #315360
CAUTERY	Megadyne needle tip (0118A), [2] holsters ✓ Safe-t-Vac, ✓ [2] holsters, ✓ Bovie 20/20 pure
SUCTION	[2] blue 3/16" suction tubing for Saf-T-Vac & Yankauer
SOLUTIONS IRRIGATION	[1] liters NS irrigation, [1] saline bowl - NO ICE unless doing FACELIFT
NEEDLES SYRINGES	[2] 27g 1" needle on [2] 10cc syringe [2] 16g angiocath [2] 30cc syringes ➤ for Gwen irrigation
LOCAL ANESTHETIC	Solution E1 1% lidocaine w/ epi - 1 vial, buffered 1:10 with HCO ₃
MEDICATIONS	[4] Versed; [4] Fentanyl; [1] Zofran; [1] Kefzol
STERILE SUPPLIES	1000 drape, needle box, magnetic pad NO scalpel blades - will use cautery for intraoral incision
SPONGES	[2] packs Raytec, 10ea.
NON-STERILE SUPPLIES	

419 Excision Buccal Fat Pads (intra-oral approach)

page 2

SETS TO OPEN	Dr. C Facelift Set, Special Plastic Set			
INSTRUMENTS USED ON MAYO STAND				
■ FORCEPS	81 and 83 6" Adsons, smooth Adson, 7" Debakey, Dressing forceps, Cushings			
■ SCISSORS	str Metz, str Mayo, Reynolds			
■ NEEDLE HOLDERS	[2] Aesculap serr NH, [1] 6" Debakey NH			
■ RETRACTORS	Weitlaner self-retainer, sharp, medium 3/8" baby Deaver, L-shaped retractor, Weider cheek			
■ KNIFE BLADES	NONE - will use needle cautery for intraoral incision			
■ OTHER	Weitlaner self-retainer, sharp, medium			
SUTURES	No	Type	SN	status
●				
●	[4]	4-0 Chromic PS-2	1637	open 2 hold 2
●	[2]	3-0 nylon PS-1	1663	open 1
●				
●				
●				
DRESSINGS	none			
POSTOPERATIVE	Remove head cushion, put 2 nd pillow under head			
DATES MODIFIED	Dec 2016, Jan 2017			

NOTE DIFFERENCES:

Solution E-1 local

No scalpel, will use short needle tip electrocautery Megadyne (0118A)

Weitlaner self-retainer, sharp, medium

Tongue stitch using 3-0 nylon