

SPLIT THICKNESS SKIN GRAFT	
Donor site usually taken from thigh or buttock. Procedure involves preparation of recipient site (face, hand, leg or foot) and harvesting of skin graft from donor site. Often done in conjunction with other reconstructive procedure, as in 4 th Stage Microtia	
OPERATIVE TIME	1 hour
ICD9 CODE:	depends on recipient site procedure
CPT CODE(S):	15100 STSG
ANESTHESIA NOTES	General
PATIENT POSITION SETUP	On table, may prop thigh or hip under folded towel
TABLE POSITION	depends on recipient site
ROOM SETUP	pre-warm room and bed, Baer hugger, multi-access
EQUIPMENT SPECIALS	Padgett electric dermatome, power box, blade Mesher and plastic mesh sheets - ask
PREP	CHG prep to donor site, thigh or buttock Technicare to face
DRAPING	[4] 1000 drapes, stickies
CAUTERY	none for the harvest
SUCTION	none for the harvest
SOLUTIONS IRRIGATION	Place 1mg epinephrine in kidney basin Cefazolin irrigation for graft
NEEDLES SYRINGES	Ask, 10cc syringe, 25g 1 ½" needle
LOCAL ANESTHETIC	Ask, may inject with local.
MEDICATIONS	<input type="checkbox"/> Epinephrine 1mg vial - draw up <input type="checkbox"/> Cefazolin for irrigation <input type="checkbox"/> Local - ask
SETS TO OPEN	Plastic Set
INSTRUMENTS USED (ON MAYO STAND)	
■ FORCEPS	smooth adson, 03, 05
■ SCISSORS	stevens, iris, metz
■ NEEDLE HOLDERS	crile wood, webster
■ RETRACTORS SKIN HOOKS	skin hooks
■ KNIFE BLADES	No. 11 blade for piercing graft
■ OTHER	Mesher - ask
STERILE SUPPLIES	Glass medicine cup, Mastisol Large Telfa
SPONGES	4x4 raytec
NON-STERILE SUPPLIES	

	No	Type	SN	status
SUTURES	[4]	5-0 Chromic P3		hold
	[2]	4-0 Vicryl P3		hold
	[]			
DRAINS	none			
DRESSINGS	<u>Recipient site:</u> ask <u>Donor site:</u> Large Tegaderm, mastisol, abd, kerlix, ace			
POSTOPERATIVE NOTES				
COMMENTS				
DATE MODIFIED	March 23, 2007			

[Back to Top](#)
[Back to Master List](#)